## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



1.9315

## INSECT PEST SURVEY

Bureau of Entomology, U. S. D. A.,

State Entomological Agencies cooperating.

Special Report No. 28 November 15, 1923.

POTATO TUBER MOTH IN VIRGINIA. (Phthorimaea operculella Zell.)

The outbreak of potato tuber moth on the Eastern Shore of Virginia extends well up into Accomac County. Careful scouting during the last two weeks has established the region of maximum damage to be between Eastville and Exmore. South of Eastville larvae of the insect were found in fewer numbers as far down the peninsula as Kiptopeke. North of Exmore the survey is still unfinished, but larvae have been taken at Onley, Onancock, Parksley and Bloxom. At the last place only one larva was found by five inspectors, who looked over many fields.

The Eastern Short of Virginia Produce Exchange, which handles practically all of the potatoes of the district, has agreed not to sell any seed from Virginia, or ship any, until January 1, 1924. By that time the condition of stored home-grown seed with respect to tuber worm infestation can be easily determined by their inspectors. No infested seed will be accepted by the exchange for shipment.

A tuber moth campaign has been planned and started in the affected region by the Virginia Truck Experiment Station. Arrangements have been made to fumigate most of the home-grown seed before planting time. To date two storage houses of a combined capacity of 250,000 cubic feet have been treated.

(Extract from letter from Herbert Spencer, of the Virginia Truck Experiment Station, November 11, 1923, Norfolk, Virginia.)

## VERRET PRET AURUST

Bureau of Entendiogy, U. D. I. A.,

Elate Emigratedical Agencies conversible

Sales at Report to the the transfer of the tra

## POTUTO TURKE FORM IN PRESIDENTS.

The outpreak of potato stoer acts on the finters in the fortant facts of Virtinia exists well up late Actorse tourity. Careful accution during the lest two weste mes established the region of rexional demans to up between heatylistand Engage. South of Eastylist leaves of the indeed were found in fewer cumbers on far down the pentionia as Kiptopeke. North of ixense the survey is still unfitudent, but large near taken at Onley, Descook, Farkeley and Blower, at the last place only one large was frue by Five last place only one large was frue by Five last potent over sony fields.

The Eastern Chart of Virginia Produce Exchange, which bardles produceing all of the potatoes of the district, has accord not to sail any send from Virginia, or ship say, until lenuary 1, 10%. By the same the constituent of stored here-erown seed till range of to there were work infinitely determined by their important. We infraged need will be accepted by the exchange for shirtent.

A tuber moth empedan has been planned and started in the alfacted region by the Virginia Track Experiment Stallon. Arrangements have been made to Iuminote most of the home-ground and before
planting time. To date two storens housen of a constant oppacity
of 250,000 costs feet have been treated;

(Entract from letter from Horoset Speacot, Of the Virginia from North Parks (Parks 13, North 13)